Call for Paper

第三屆亞太醫用超音波新進展國際論壇 、
暨中華民國醫用超音波學會 37 週年暨 2021 年年會聯合會
3rd Asia Pacific international Symposium on Advances in Medical Ultrasound、
37th Anniversary & 2021 Annual Convention of Taiwan Society of Ultrasound in Medicine

October 16-17, 2021

地點:National Taiwan University College of Medicine

General Information

Abstracts should include background, materials and methods, results and conclusion. Do not include references or acknowledgements. The length of the abstract should not exceed 300 words, no figures. All abstracts must be written in English.

Title: The title should be first letter capital.
Authors: Type names of authors, institution, city and country.
Key words: Not more than three to five key words or short phrases.

On-line Submission: 請您至學會網站 www.sumroc.org.tw 點選年會專區 - 線上投稿。

Deadline: July 31, 2021

Sample：

Biometric Difference in Primary Angle-Closure Glaucoma: Study on Lens

Chong-Bin Tsai and Por-Tying Hung1
Department of Ophthalmology, Chiayi Christian Hospital, Chiayi, and 1Department of Ophthalmology, National Taiwan University Hospital, Taipei

Background: As a couse of shallow anterior chamber, certain variables of the lens are considered to be important risk factors for primary angle-closure glaucoma.
Materials and Methods: Using A-scan ultrasound, intraindividual comparisons of eye lens thickness were carried out in 41 patients with mature cataract in one eye and intumescent lens in the other.
Results: The average thickness of an intumescent lens (4.52±0.50mm) is greater than that of a mature lens (4.02±0.62mm), ($p<0.001$). No significant difference existed in the depth of the anterior chamber or axial length.
Conclusion: The A-scan results confirmed the importance of lens factors in primary angle-closure glaucoma involving "constitutional" or hereditary elements, as well as lens growth form aging and intumescent lens during cataract formation.
(**Key words:** A-scan ultrasonography, primary angle-closure glaucoma, intumescent lens, cataractous lens)