

Call for Paper

第七屆亞太醫用超音波新進展國際論壇、
中華民國醫用超音波學會

41週年暨2025年年會暨第一次學術研討會

7th Asia Pacific international Symposium on Advances in Medical Ultrasound
and

41th Anniversary & 2025 Annual Convention of Taiwan Society of Ultrasound in Medicine

October 18-19, 2025

地點：臺大醫學院基醫學大樓

General Information

Abstracts should include background, materials and methods, results and conclusion. Do not include references or acknowledgements. The length of the abstract should not exceed 300 words, no figures. All abstracts must be written in English.

Title: The title should be first letter capital.

Authors: Type names of authors, institution, city and country.

Key words: Not more than three to five key words or short phrases.

On-line Submission: 請您至學會網站 www.sumroc.org.tw 點選年會專區 - 線上投稿。

Submission Deadline: July 31, 2025

Sample :

Biometric Difference in Primary Angle-Closure Glaucoma: Study on Lens

Chong-Bin Tsai¹ and Por-Tying Hung²

¹Department of Ophthalmology, Chiayi Christian Hospital, Chiayi, and ²Department of Ophthalmology, National Taiwan University Hospital, Taipei

Background: As a cause of shallow anterior chamber, certain variables of the lens are considered to be important risk factors for primary angle-closure glaucoma.

Materials and Methods: Using A-scan ultrasound, intraindividual comparisons of eye lens thickness were carried out in 41 patients with mature cataract in one eye and intumescent lens in the other.

Results: The average thickness of an intumescent lens (4.52 ± 0.50 mm) is greater than that of a mature lens (4.02 ± 0.62 mm), ($p < 0.001$). No significant difference existed in the depth of the anterior chamber or axial length.

Conclusion: The A-scan results confirmed the importance of lens factors in primary angle-closure glaucoma involving "constitutional" or hereditary elements, as well as lens growth form aging and intumescent lens during cataract formation.

(Key words: A-scan ultrasonography, primary angle-closure glaucoma, intumescent lens, cataractous lens)